



Original Research Article

Aloe vera: Botanical Characteristics, Phytochemical Properties, Therapeutic Potential and Biotechnological Applications

Sarla Rani^{1*} and Mahesh Kumar²

¹Pt. C.L.S. Government College, Karnal, Haryana, India

²IIHS, Kurukshetra University, Kurukshetra, Haryana, India

Article Information: Received on 25-10-2021 Accepted on 27-12-2021 Available online 29/12 /2021

Abstract

Aloe vera is an important medicinal plant that has been used traditionally for several centuries and has subsequently attracted considerable scientific and industrial attention. The plant is particularly recognized for its succulent leaves, which contain two chemically distinct materials: the inner clear gel and the yellowish latex located beneath the leaf rind. These fractions differ in composition and consequently possess different biological and safety profiles. Aloe gel contains polysaccharides, amino acids, minerals, vitamins, enzymes, phenolic compounds and other constituents, whereas the latex contains anthraquinone derivatives such as aloin. Acemannan and other polysaccharides have been investigated for their possible roles in immunological responses, cellular activity and tissue repair. Experimental studies have also reported antioxidant, anti-inflammatory and antimicrobial activities associated with different aloe constituents. Aloe has received particular attention in relation to wound healing, dermatological applications and metabolic health. However, clinical evidence is variable, and differences in plant material, extraction procedures, dosage and product composition make comparison among studies difficult. Safety is especially important for whole-leaf and latex-containing preparations because anthraquinone-associated constituents may produce adverse effects. Beyond therapeutic applications, Aloe vera has considerable importance in the food, cosmetic, pharmaceutical and biotechnology sectors. Tissue culture, micropagation, analytical characterization, standardized extraction and development of aloe-derived biomaterials represent promising areas for future research. This review summarizes the botanical features, major phytochemical constituents, biological properties, health-related applications, safety considerations, industrial significance and future research opportunities associated with Aloe vera.

Keywords: Aloe Vera; Medicinal Plant; Phytochemicals; Acemannan; Anthraquinones; Antioxidant Activity; Wound Healing; Biotechnology; Medicinal Applications; Safety

Introduction

Medicinal plants have played a significant role in traditional healthcare and continue to provide important sources of biologically active compounds for modern research. Among these plants, Aloe vera has received extensive scientific attention because of its historical medicinal use and broad application in pharmaceutical, cosmetic, food and nutraceutical products.

The plant is valued particularly for its fleshy leaves, which contain an inner transparent gel and a yellowish latex fraction. Although both materials originate from the same leaf, their chemical composition and biological properties are substantially different [1]-[3].

Traditional applications of Aloe vera include its use for minor skin injuries, burns, dermatological problems and gastrointestinal complaints. Modern investigations have examined a wider range of biological properties, including antioxidant, antimicrobial, anti-inflammatory, immunomodulatory and wound-related activities [4], [5]. Nevertheless, the presence of biological activity in experimental systems does not necessarily establish clinical effectiveness in humans. Reviews of the clinical literature have identified

*Corresponding author: **Sarla Rani**

DOI: <http://doi.org/10.54618/IJMAHS.2021125>

This is an open-access article, which permits the use and distribution of article provided that original author and source are credited

considerable variation in the strength and quality of evidence for different applications [6].

Interest in Aloe vera has expanded beyond traditional medicine. Aloe-derived preparations are now incorporated into cosmetic formulations, beverages, functional foods, topical products and other commercial preparations [2], [7]. Consequently, understanding the plant at botanical, chemical, biological and technological levels is essential for the development of safe and standardized products.

Botanical Characteristics of Aloe vera

Aloe vera is a perennial succulent adapted to environments where water availability may be limited. Its fleshy leaves possess water-storage tissues that contribute to its ability to withstand relatively dry conditions. The leaves are generally arranged in a compact rosette and range from green to greyish-green, with characteristic marginal spines [1].

The plant commonly possesses a short stem from which numerous thick leaves emerge. Vegetative multiplication can occur through offsets or suckers, an attribute that facilitates cultivation and propagation. The flowering stalk produces tubular yellowish flowers arranged in racemes [4]. The leaf can broadly be divided into three structural regions: the external rind, the vascular region associated with the bitter latex, and the inner parenchymatous tissue containing the clear gel. This anatomical organization is scientifically important because the chemical composition and biological activity of an aloe preparation depend strongly on the specific leaf fraction used [3].

Aloe Gel and Aloe Latex

The inner gel represents the principal water-rich component of the leaf. In addition to water, it contains polysaccharides, amino acids, minerals, vitamins, enzymes and several other compounds. Among the polysaccharides, acemannan and related mannose-rich polymers have been extensively studied because of their possible involvement in immune responses and tissue repair [3], [5]. Aloe latex is chemically distinct from the inner gel. It is a yellowish and bitter exudate located close to the leaf rind and contains anthraquinone derivatives, including aloin. These compounds are associated with the traditional laxative properties of aloe [8].

The distinction between gel and latex is therefore essential in both research and clinical interpretation. Products described as inner-leaf gel, decolorized whole-leaf extract, whole-leaf preparation or latex should not automatically be considered equivalent because their chemical composition and toxicological characteristics may differ considerably [9].

Phytochemical Constituents

The biological properties of Aloe vera are associated with a chemically diverse collection of compounds. The

composition of aloe preparations is influenced by plant material, cultivation conditions, harvesting, processing and storage. Consequently, commercially available preparations may vary considerably in their chemical profiles [5].

1. Polysaccharides

Polysaccharides are among the most extensively investigated constituents of aloe gel. Acemannan, an acetylated mannose-rich polysaccharide, has received particular attention. Experimental research has explored its potential effects on immune-related responses, cellular processes and tissue repair [3]. The molecular characteristics of aloe polysaccharides may change during processing and storage. Such alterations can influence the biological properties of the final preparation and may partly explain differences observed between aloe products [3], [10].

2. Anthraquinones

Anthraquinone derivatives are principally associated with the latex fraction. Aloin is among the best-known compounds in this group and is responsible for the laxative activity historically attributed to aloe latex [8]. At the same time, anthraquinones are important from a safety perspective. Toxicological concerns have been reported for certain whole-leaf or anthraquinone-containing preparations, making the controlled removal or reduction of these compounds an important aspect of aloe processing [9].

3. Phenolic and Chromone Compounds

Aloe contains phenolic substances and chromone derivatives that contribute to its chemical diversity. Experimental investigations have associated some of these compounds with antioxidant and other biological activities [1], [5].

4. Flavonoids

Flavonoid and related phenolic compounds have been detected in aloe preparations. Their potential antioxidant and anti-inflammatory activities have contributed to continuing interest in the pharmacological properties of the plant [5].

5. Sterols

Phytosterols and other lipid-soluble compounds are also reported constituents of Aloe vera. Experimental research has investigated their possible relationship with inflammatory and metabolic processes [3].

6. Vitamins, Minerals and Amino Acids

Aloe gel contains a variety of vitamins, minerals and amino acids. However, the concentrations of these substances are influenced by cultivar, environmental

conditions, cultivation practices and processing. Therefore, the mere detection of these nutrients should not be interpreted as evidence of a clinically significant nutritional effect [2].

Biological and Pharmacological Properties

1. Antioxidant Activity

Oxidative stress results from an imbalance involving reactive oxygen species and the body's protective mechanisms. Several constituents present in Aloe vera extracts have demonstrated antioxidant activity under experimental conditions. Phenolic compounds, flavonoids and other phytochemicals may contribute to this activity [1], [5].

These findings provide a scientific rationale for continued investigation of aloe in conditions associated with oxidative stress. However, antioxidant activity demonstrated in laboratory assays should not be directly interpreted as proof that oral aloe consumption prevents human disease. Clinical effectiveness requires appropriately designed human studies [6].

2. Anti-inflammatory Activity

Inflammatory responses participate in numerous physiological and pathological processes. Various constituents of Aloe vera have been investigated for their possible effects on inflammatory mediators and cellular signaling pathways [5]. Experimental observations provide a possible explanation for some of the traditional topical applications of aloe. Nevertheless, findings from cell and animal studies should be carefully distinguished from evidence generated through controlled human clinical research.

3. Antimicrobial Activity

Laboratory investigations have demonstrated antimicrobial effects from selected aloe preparations against certain microorganisms. The magnitude of this activity may depend upon the concentration of the extract, extraction method, chemical composition and microorganism under investigation [5]. Although these findings are scientifically relevant, aloe preparations should not automatically be considered substitutes for clinically established antimicrobial treatments.

Wound-Healing Potential

Wound repair is one of the most frequently investigated applications of Aloe vera. Experimental research indicates that aloe constituents may influence inflammatory responses, cellular proliferation, collagen-associated processes and tissue repair [11], [12].

A systematic review of research on cutaneous wounds reported indications of potentially beneficial effects while also emphasizing limitations in the

available evidence and the need for additional high-quality studies [11]. Earlier clinical evidence concerning burn wounds also suggested shorter healing times in some aloe-treated groups, although differences among products and study methodologies prevented definitive conclusions [13].

Evidence should therefore be interpreted cautiously. A Cochrane review concluded that available trials involving acute and chronic wounds were generally small and methodologically limited, and that high-quality evidence was insufficient to establish aloe as a definitive wound treatment [14]. Thus, although wound-healing research remains promising, further well-designed clinical studies are required.

Potential Effects on Diabetes and Blood Glucose

The possible relationship between Aloe vera consumption and glucose regulation has been investigated in people with prediabetes and type 2 diabetes. A systematic review and meta-analysis by Suksomboon et al. [15] reported evidence suggesting possible improvements in certain measures of glycemic control following aloe supplementation, although substantial variation existed among the studies.

Potential changes have also been reported for fasting blood glucose and HbA1c, together with some lipid parameters. However, limitations in the number and quality of randomized trials restrict the strength of the conclusions [15]. An overview of systematic reviews similarly identified potentially favorable effects on selected glycemic outcomes but emphasized differences between preparations and study designs [16]. Accordingly, aloe remains an area of research in metabolic health rather than an established replacement for conventional diabetes management.

Effects on Blood Lipids

The influence of aloe preparations on lipid metabolism has also been examined. Earlier clinical research suggested that oral aloe preparations might influence blood lipid concentrations in individuals with hyperlipidemia [17]. However, differences in preparation, dose, duration of administration and study methodology make it difficult to define a uniform therapeutic effect. Further controlled clinical investigations are required before firm conclusions regarding the clinical significance of these observations can be drawn.

Dermatological Applications

Aloe vera gel is widely incorporated into topical and cosmetic formulations. Its high-water content and polysaccharide-rich composition make it a useful botanical ingredient in moisturizing and skin-care preparations [3].

Research has examined aloe in relation to several dermatological conditions, including burns, psoriasis and other skin disorders. Clinical evidence has

suggested possible benefits for selected applications, but findings have not been sufficiently consistent across all conditions [17]. The effectiveness of aloe therefore appears to depend upon the condition being treated, the preparation used and the method of application.

Gastrointestinal Applications

The traditional gastrointestinal use of aloe is primarily associated with its latex rather than the purified inner gel. Anthraquinone compounds such as aloin can stimulate intestinal activity and produce a laxative effect [8].

However, prolonged or excessive exposure to latex-containing preparations may result in undesirable gastrointestinal effects and disturbances in electrolyte balance. Toxicological concerns associated with whole-leaf and latex preparations further emphasize the importance of distinguishing between different aloe products [9]. Consequently, traditional use of aloe should not be interpreted as evidence that all forms of aloe are appropriate for oral gastrointestinal applications.

Safety and Toxicological Considerations

Safety is a fundamental consideration when evaluating medicinal plants and their commercial preparations. In the case of Aloe vera, safety cannot be considered independently of the specific portion of the plant and processing procedure used.

Aloe gel, whole-leaf extract and latex possess different chemical profiles. Oral exposure to certain aloe preparations has been associated with adverse effects, including gastrointestinal disturbances and electrolyte abnormalities. Toxicological concerns have also been raised regarding preparations containing anthraquinone-related compounds [9]. For this reason, scientific studies should clearly identify the type of aloe preparation investigated. Standardized processing, appropriate quality control and chemical characterization are particularly important for products intended for oral consumption.

Food, Cosmetic and Industrial Applications

The expanding commercial importance of Aloe vera is reflected in its incorporation into food, cosmetic and pharmaceutical products. Aloe gel has been investigated as a functional food ingredient because of its polysaccharide content and other bioactive constituents [2], [7].

Aloe-based beverages and other food products are commercially available in various markets. However, careful processing is essential because undesirable latex and anthraquinone-associated constituents may remain if the raw material is inadequately processed [7]. The cosmetic industry also uses aloe in creams, lotions, shampoos, soaps and other formulations. Its water-rich gel and established consumer acceptance

contribute to its continuing use as a botanical ingredient [3].

Biotechnological Importance

Aloe vera offers several opportunities for biotechnology research and industrial development. Important areas include tissue culture, micropropagation, extraction and purification of polysaccharides, development of standardized herbal preparations and investigation of aloe-derived biomaterials.

Plant tissue culture can facilitate the rapid multiplication of selected planting material and may assist in maintaining genetically uniform stocks. Advanced analytical techniques, including chromatographic and metabolomic approaches, can support the identification, characterization and standardization of bioactive constituents. Standardization is particularly important because cultivation, harvesting, processing and storage conditions may alter the chemical composition of aloe products. Greater chemical consistency would improve the reproducibility of experimental findings and facilitate the development of better-defined botanical preparations [3], [5].

Current Research Challenges and Future Perspectives

Despite extensive research on Aloe vera, several scientific challenges remain. One of the most important is the lack of uniformity among preparations used in different studies. Investigations may involve fresh gel, dried gel, whole-leaf extracts, decolorized extracts or latex. These materials can differ substantially in their concentrations of both biologically active and potentially undesirable compounds [9].

Another major limitation is the relatively restricted availability of large, well-controlled human clinical studies. Laboratory and animal experiments have generated numerous hypotheses regarding possible mechanisms of action, but translation of these observations into consistent clinical outcomes remains incomplete [6].

Future research should therefore concentrate on:

1. Development of standardized extraction procedures.
2. Detailed chemical characterization of aloe preparations.
3. Identification and validation of biologically active constituents.
4. Dose-response investigations.
5. Long-term safety assessment.
6. Well-designed randomized controlled clinical trials.
7. Improved quality-control systems for commercial products.

8. Development of standardized pharmaceutical and nutraceutical preparations.
9. Exploration of aloe-derived biomaterials.
10. Application of modern biotechnology and analytical technologies to aloe research.

Greater integration of pharmacognosy, biotechnology, analytical chemistry and clinical research could improve understanding of the relationship between aloe composition and biological activity.

Conclusion

Aloe vera is an important medicinal plant with a long history of traditional utilization and a substantial contemporary role in pharmaceutical, cosmetic, food and nutraceutical industries. Its biological importance is associated with a complex mixture of polysaccharides, anthraquinones, phenolic compounds, flavonoids, sterols, amino acids, enzymes and other constituents [1], [3], [5].

Experimental research supports continued investigation of aloe for antioxidant, antimicrobial, anti-inflammatory and wound-related activities. Clinical studies have additionally explored its possible influence on glycemic control and lipid metabolism, although the strength and consistency of the evidence vary considerably among applications [15], [16].

An important scientific consideration is the distinction between the inner gel and the latex fraction. Latex contains anthraquinone-associated constituents and therefore presents different pharmacological and toxicological considerations from purified inner-leaf gel [8], [9]. Overall, Aloe vera remains a valuable subject for medicinal-plant research, biotechnology, pharmacognosy and pharmaceutical science. Future progress will depend on standardized preparations, rigorous chemical characterization, appropriate safety assessment and well-designed clinical investigations. Such approaches can help distinguish traditionally attributed benefits from applications supported by reproducible scientific evidence.

References

- [1] T. Reynolds and A. C. Dweck, "Aloe vera leaf gel: A review update," *J. Ethnopharmacol.*, vol. 68, pp. 3–37, 1999.
- [2] K. Eshun and Q. He, "Aloe vera: A valuable ingredient for the food, pharmaceutical and cosmetic industries—A review," *Crit. Rev. Food Sci. Nutr.*, vol. 44, pp. 91–96, 2004.
- [3] J. H. Hamman, "Composition and applications of Aloe vera leaf gel," *Molecules*, vol. 13, pp. 1599–1616, 2008.
- [4] A. Surjushe, R. Vasani, and D. G. Saple, "Aloe vera: A short review," *Indian J. Dermatol.*, vol. 53, no. 4, pp. 163–166, 2008.
- [5] M. H. Radha and N. P. Laxmipriya, "Evaluation of biological properties and clinical effectiveness of Aloe vera: A systematic review," *J. Tradit. Complement. Med.*, vol. 5, no. 1, pp. 21–26, 2015.
- [6] S. Sadoyu et al., "Aloe vera and health outcomes: An umbrella review of systematic reviews and meta-analyses," *Phytother. Res.*, 2021.
- [7] E. R. Rodríguez, J. D. Martín, and C. D. Romero, "Aloe vera as a functional ingredient in foods," *Crit. Rev. Food Sci. Nutr.*, vol. 50, pp. 305–326, 2010.
- [8] M. D. Boudreau and F. A. Beland, "An evaluation of the biological and toxicological properties of Aloe barbadensis (Miller), Aloe vera," *J. Environ. Sci. Health C*, vol. 24, pp. 103–154, 2006.
- [9] X. Guo and N. Mei, "Aloe vera: A review of toxicity and adverse clinical effects," *J. Environ. Sci. Health C*, vol. 34, pp. 77–96, 2016.
- [10] A. Femenia, E. S. Sánchez, S. Simal, and C. Rosselló, "Compositional features of polysaccharides from Aloe vera (Aloe barbadensis Miller) plant tissues," *Carbohydr. Polym.*, vol. 39, pp. 109–117, 1999.
- [11] D. Hekmatpou, F. Mehrabi, K. Rahzani, and A. Aminiyan, "The review on properties of Aloe vera in healing of cutaneous wounds," *BioMed Res. Int.*, vol. 2015, Art. no. 714216, 2015.
- [12] D. Hekmatpou, F. Mehrabi, K. Rahzani, and A. Aminiyan, "The effect of Aloe vera clinical trials on prevention and healing of skin wound: A systematic review," *Iran. J. Med. Sci.*, vol. 44, no. 1, pp. 1–9, 2019.
- [13] R. Maenthaisong, N. Chaiyakunapruk, S. Niruntraporn, and C. Kongkaew, "The efficacy of Aloe vera used for burn wound healing: A systematic review," *Burns*, vol. 33, no. 6, pp. 713–718, 2007.
- [14] A. D. Dat, F. Poon, K. B. Pham, and J. Doust, "Aloe vera for treating acute and chronic wounds," *Cochrane Database Syst. Rev.*, 2012.
- [15] N. Suksomboon, N. Poolsup, and S. Punthanitisarn, "Effect of Aloe vera on glycaemic control in prediabetes and type 2 diabetes: A systematic review and meta-analysis," *J. Clin. Pharm. Ther.*, vol. 41, pp. 180–188, 2016.
- [16] F. Araya-Quintanilla et al., "Effectiveness of Aloe vera in patients with type 2 diabetes mellitus and pre-diabetes: An overview of systematic reviews," *Diabetes Metab. Syndr., Clin. Res. Rev.*, vol. 15, no. 6, Art. no. 102292, 2021.
- [17] B. K. Vogler and E. Ernst, "Aloe vera: A systematic review of its clinical effectiveness," *Br. J. Gen. Pract.*, vol. 49, pp. 823–828, 1999.